


This article was downloaded by: [USDA Natl Agricultul Lib]

On: 18 November 2009

Access details: Access Details: [subscription number 731827198]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Soil Science and Plant Analysis

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597241>

Role of Cover Crops in Improving Soil and Row Crop Productivity

N. K. Fageria ^a; V. C. Baligar ^b; B. A. Bailey ^b

^a National Rice and Bean Research Center of Embrapa, Santo Antônio de Goiás, Goiás, Brazil ^b USDA-ARS-Alternate Crops and Systems Research Laboratory, Beltsville Agricultural Research Center, Beltsville, Maryland, USA

To cite this Article Fageria, N. K., Baligar, V. C. and Bailey, B. A. 'Role of Cover Crops in Improving Soil and Row Crop Productivity', Communications in Soil Science and Plant Analysis, 36: 19, 2733 — 2757

To link to this Article: DOI: 10.1080/00103620500303939

URL: <http://dx.doi.org/10.1080/00103620500303939>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Role of Cover Crops in Improving Soil and Row Crop Productivity

N. K. Fageria

National Rice and Bean Research Center of Embrapa,
Santo Antônio de Goiás, Goiás, Brazil

V. C. Baligar and B. A. Bailey

USDA-ARS-Alternate Crops and Systems Research Laboratory,
Beltsville Agricultural Research Center, Beltsville, Maryland, USA

Abstract: Cover crops play an important role in improving productivity of subsequent row crops by improving soil physical, chemical, and biological properties. The objective of this article is to review recent advances in cover crops practice, in the context of potential benefits and drawbacks for annual crop production and sustained soil quality. Desirable attributes of a cover crop are the ability to establish rapidly under less than ideal conditions, provide sufficient dry matter or soil cover, fix atmospheric nitrogen (N), establish a deep root system to facilitate nutrient uptake from lower soil depths, produce organic matter with low-residue carbon/nitrogen (C/N) ratio, and absence of phytotoxic or allelopathic effects on subsequent crops. Cover crops can be leguminous or nonleguminous. Leguminous cover crops provide a substantial amount of biologically fixed N to the primary crop, as well as ease of decomposition due to their low C/N ratio. Legume cover crops also possess a strong ability to absorb low available nutrients in the soil profile and can help in increasing concentration of plant nutrients in the surface layers of soil. Some nonleguminous cover crops having high N scavenger capacity compared with leguminous crops and sometimes, the growth of these scavenging grass cover crops is limited by N deficiency, growing grass/legume mixtures appears to be the best strategy in obtaining maximum benefits from cover crops.

Keywords: Biological activities, organic matter, soil quality, sustainable system

Received 27 August 2004, Accepted 14 July 2005

Address correspondence to N. K. Fageria, National Rice and Bean Research Center of Embrapa, Caixa Postal 179, Santo Antônio de Goiás, Goiás, CEP 75375-000, Brazil.
E-mail: fageria@cnpaf.embrapa.br

INTRODUCTION

Increasing crop productivity and maintaining a clean environment are major challenges to agricultural scientists in the 21st century. To meet these challenges, crop production practices need to be modified in favor of higher yields and minimized environmental pollution. The management of crop residues is a key component of sustainable cropping systems (Ruffo and Bollero, 2003). Historically, crop residues have played an important role as mulch for soil and water conservation and as an input for maintaining soil organic matter and returning nutrients to soil. To achieve these objectives the use of cover crops in cropping systems is an important strategy. Before discussing cover crops and their role in crop production, it is important to define the term cover crops. Cover crops can be defined as close-growing crops that provide soil protection, and soil improvement between periods of normal crop production, or between trees in orchards and vines in vineyards (SSSA, 1997). Cover crops are grown not for market purposes, but when plowed under and incorporated into the soil, cover crops may be referred to as green manure crops. Cover crops are sometimes called catch crops. In Table 1 are listed important tropical and temperate cover crops. Cover crops are usually killed on the soil surface before they are mature by using appropriate herbicides. In most studies, cover crops managed as no-till mulches have been killed with glyphosate (N-[phosphono-methyl]glycine), paraquat (1,1-dimethyl-4-bipyridinium ion) (Bauer and Reeves, 1999; Raimbault et al., 1990; Reeves et al., 1993; Sarrantonio and Scoot, 1988) or mixture of non-selective, post-emergence and pre-emergence herbicide (Teasdale and Shirley, 1998). Because many growers often want to reduce use of chemical inputs, non-chemical methods of killing or suppressing cover crops are desirable (Creamer and Dabney, 2002). These mechanical methods are mowing, rolling, roll-chopping, undercutting, and partial rototilling. Creamer and Dabney (2002) have reviewed literature on killing cover crops mechanically.

Cover crops are generally included in cropping systems as nutrient management tools (Ruffo and Bollero, 2003). Cover crops can be leguminous or nonleguminous. Legume cover crops are used as a source of nitrogen (N) for the following cash crop (Smith et al., 1987) while grasses are mainly used to reduce NO₃ leaching and erosion (Meisinger et al., 1991). Biological N fixation by leguminous crops offers potential to reduce the need for N fertilizers for the succeeding crop (Singh et al., 2004). A bicultural of a legume and a grass is used with the intention of providing both benefits simultaneously (Ranells and Waggoner, 1996).

The benefit from legume cover crops in crop rotation has long been recognized and believed to be mainly attributed to N contribution to subsequent crops (Smith et al., 1987). In recent years, economic and environmental considerations have renewed interest in this old practice for improving crop productivity and soil health and maintaining sustainability of agroecosystems.

Table 1. Major cover crops of tropical and temperate regions

Tropical region		Temperate region	
Common name	Scientific name	Common name	Scientific name
Sunnhemp	<i>Crotalaria juncea</i> L.	Hairy vetch	<i>Vicia villosa</i> Roth
Sesbania	<i>Sesbania aculeata</i> Retz Poir	Barrel medic	<i>Medicago truncatula</i> Gaertn
Sesbania	<i>Sesbania rostrata</i> Bremek & Oberm	Alfafa	<i>Medicago sativa</i> L.
Cowpea	<i>Vigna unguiculata</i> L. Walp.	Black lentil	<i>Lens culinaris</i> Medikus
Soybean	<i>Glycine max</i> L. Merr.	Red clover	<i>Trifolium pratense</i> L.
Clusterbean	<i>Cyamopsis tetragonoloba</i>	Soybean	<i>Glycine max</i> L. Merr.
Alfalfa	<i>Medicago sativa</i> L.	Faba bean	<i>Vicia faba</i> L.
Egyptian clover	<i>Trifolium alexandrium</i> L.	Crimson clover	<i>Trifolium incarnatum</i> L.
Wild indigo	<i>Indigofera tinctoria</i> L.	Ladino clover	<i>Trifolium repens</i> L.
Pigeon pea	<i>Cajanus cajan</i> L. Millspaugh	Subterranean clover	<i>Trifolium subterraneum</i> L.
Mungbean	<i>Vigna radiata</i> L. Wilczek	Common vetch	<i>Vicia sativa</i> L.
Lablab	<i>Lablab purpureus</i> L.	Purple vetch	<i>Vicia benghalensis</i> L.
Graybean	<i>Mucuna cinerecun</i> L.	Cura clover	<i>Trifolium ambiguum</i> Bieb.
Buffalobean	<i>Mucuna aterrima</i> L. Piper & Tracy	Sweet clover	<i>Melilotus officinalis</i> L.
Crotalaria		Winter pea	<i>Pisum sativum</i> L.
Breviflora	<i>Crotalaria breviflora</i>	Narrowleaf vetch	<i>Vicia angustifolia</i> L.
White lupin	<i>Lupinus albus</i> L.	Milk vetch	<i>Artragalus sinicus</i> L.
Milk vetch	<i>Astragalus sinicus</i> L.		
Crotalaria	<i>Crotalaria striata</i>		
Zornia	<i>Zornia latifolia</i>		
Jackbean	<i>Canavalia ensiformis</i> L. DC.		
Tropical kudzu	<i>Pueraria phaseoloides</i> (Roxb.) Benth.		
Velvetbean	<i>Mucuna deeringiana</i> Bort. Merr.		
Adzuki bean	<i>Vigna angularis</i>		
Brazilian stylo	<i>Stylosanthes guianensis</i>		
Jumbiebean	<i>Leucaena leucocephala</i> Lam. De Wit		
Desmodium	<i>Desmodiumovalifolium</i> Guillemin & Perrottet		
Pueraria	<i>Pueraria phaseoloides</i> Roxb.		

The objective of this review is to compile information on beneficial effects and drawbacks of cover crops on soil health, and crop yield.

BENEFITS OF COVER CROPS

Planting cover crops before or between main crops as well as between trees or shrubs of plantation crops can improve soil physical, chemical, and biological properties and consequently lead to improved soil health and yield of principal crops. Leaving cover crops as surface mulches in no-till crop production systems has the advantage of increasing nitrogen economy (Smith et al., 1987; Frye et al., 1988; Sustainable Agriculture Network, 1998) conserving soil moisture (Morse, 1993), reducing soil erosion (Langdale et al., 1991), improving soil physical properties (Blevins and Frye, 1993), increasing nutrient retention (Staver and Brinsfield, 1998; Dinnes et al., 2002), increasing soil fertility (Cavigelli and Thien, 2003), suppressing weeds (Creamer and Baldwin, 2000; Creamer et al., 1996a), reducing diseases and insects (Sustainable Agriculture Network, 1998; Ristaino et al., 1996), reducing global warming potential (Robertson et al., 2000), and increasing crop yields (Triplett et al., 1996). These various beneficial effects of cover crops on soil management and crop productivity are discussed in the following sections.

Nitrogen Economy

The contribution of N is the most commonly observed primary benefit of leguminous crops (Singh et al., 1992). Both legume and nonlegume cover crops affect N fertilizer management (Bauer and Roof, 2004). Legume cover crops fix atmospheric N and reduce N fertilizer needs for succeeding cash crops (Hoyt and Hargrove, 1986; Reeves, 1994). The rate of N fixed by cover crops is determined largely by the genetic potential of the legume species and by the amount of plant available N in the soil. Data in Table 2 show quantities of nitrogen fixed by different legume cover crops. Two bacterial species, i.e., *Rhizobium* and *Bradyrhizobium* are responsible for symbiotic nitrogen fixation in legumes. The genus *Rhizobium* contains fast growing, acid-producing bacteria, while the *Bradyrhizobium* are slow growers that do not produce acid (Brady and Weil, 2002). Soil factors such as pH, moisture content, and temperature also determine N fixation capacity of a legume cover crop. In some cases, the amount of N provided by legume cover crops is adequate to produce optimal yields of subsequent non-leguminous crops; however, higher N requiring cereals such as corn (*Zea mays* L.) generally need supplemental N fertilizer. In such crops N fertilizer rates could be lowered appreciably while maintaining optimal economic yields (Frye et al., 1988). Table 3 shows data relating dry matter production and N uptake by different cover crops. Dry matter yields and N uptake values

Table 2. Quantity of nitrogen fixed by legume cover crops

Crop species	N ₂ Fixed (kg ha ⁻¹ crop ⁻¹)	References
Peanut (<i>Arachis hypogaea</i> L.)	40–80	Brady and Weil (2002)
Cowpea (<i>Vigna unguiculata</i> L. Walp.)	30–50	Brady and Weil (2002)
Alfalfa (<i>Medicago sativa</i> L.)	78–222	Heichel (1987)
Soybean (<i>Glycine max</i> L.)	50–150	Brady and Weil (2002)
Fava bean (<i>Vicia faba</i> L.)	177–250	Heichel (1987)
Hairy vetch (<i>Vicia villosa</i> Roth.)	50–100	Brady and Weil (2002)
Ladino clover (<i>Trifolium repens</i> L.)	164–187	Heichel (1987)
Red clover (<i>Trifolium pratense</i> L.)	68–113	Heichel (1987)
White lupine (<i>Lupinus albus</i> L.)	50–100	Brady and Weil (2002)
Field peas (<i>Pisum sativum</i> L.)	174–195	Heichel (1987)
Chickpea (<i>Cicer arietinum</i> L.)	24–84	Heichel (1987)
Pigeon pea (<i>Cajanus cajan</i> L. Huth.)	150–280	Brady and Weil (2002)
Kudzu (<i>Pueraria phaseoloides</i> Roxb. Benth)	100–140	Brady and Weil (2002)
Chick pea (<i>Cicer arietinum</i> L.)	24–84	Heichel (1987)
Greengram (<i>Vigna radiata</i> L. Wilczek.)	71–112	Chapman and Myers (1987)
Lentil (<i>Lens culinaris</i> L.)	57–111	Smith et al. (1987)

provide an indication of N uptake capacity of a cover crop from residual soil inorganic N and mineralized soil organic N as well as biological fixed N in the case of a legume cover crop. Data in Table 3 show that most legume crops have a high capacity to accumulate N in their dry matter. Similarly, rye (*Secale cereale* L.) as a cereal also accumulated 100 kg N ha⁻¹. Table 4 provides data of nitrogen fertilizer equivalence (NFE) of legume cover

Table 3. Dry matter yield and N uptake by cover crops

Crop species	Dry matter yield (Mg ha ⁻¹)	N uptake (kg ha ⁻¹)
Hairy vetch	5.1	209
Bigflower vetch	1.9	60
Crimson clover	2.4	56
Berseem clover	1.5	45
Australian winter pea	1.6	68
Common vetch	4.3	134
Subterranean clover	4.0	114
Rye	6.3	100
Wheat	1.5	29

Source: Compiled from Frye et al. (1988).

Table 4. Nitrogen fertilizer equivalence (NFE) of legume cover crops to succeeding nonlegume crops

Legume/non-legume crop	NFE (kg ha ⁻¹)
Hairy vetch/cotton	67–101
Hairy vetch + rye/corn	56–112
Hairy vetch/corn	78
Hairy vetch/sorghum	89
Hairy vetch/corn	78
Hairy vetch + wheat/corn	56
Crimson clover/cotton	34–67
Crimson clover/corn	50
Crimson clover/sorghum	19–128
Common vetch/sorghum	30–83
Bigflower vetch/corn	50
Subterranean clover/sorghum	12–103
Sesbani/allowland rice	50
Alfalfa/corn	62
Alfalfa/wheat	20–70
Arachis spp/wheat	28
Subterranean clover/wheat	66
White lupin/wheat	22–182
Arachis spp/corn	60
Pigeon pea/corn	38–49
Sesbania/potato	48
Mungbean/potato	34–148
Chickpea/wheat	15–65

Source: Compiled from Smith et al. (1987) and Kumar and Goh (2000).

crops to succeeding non-legume crops. The NFE values varied from 12 to 182 kg ha⁻¹. Smith et al. (1987) reported that NFE values range from 40 to 200 kg ha⁻¹, but more typically are between 75 to 100 kg ha⁻¹. Interseeding red clover (*Trifolium pretense* L.) into small grains is a common practice in the northeastern United States (Singer and Cox, 1998), and such practice can provide up to 85 kg N ha⁻¹ to the subsequent corn crop (Vyn et al., 1999). Researchers in the southeastern United States have estimated that legumes such as hairy vetch can supply well over 100 kg N ha⁻¹ to following corn or grain sorghum crops (Blevins et al., 1990; Hargrove, 1986; Oyer and Touchton, 1990; Waggar, 1989). On prairie soils in Kansas, Sweeney and Moyer (2004) found that grain sorghum following initial kill-down of red clover and hairy vetch yielded as much as 131% more than continuous sorghum with estimated fertilizer N equivalencies exceeding 135 kg ha⁻¹.

Legume cover crops should be inoculated with an appropriate strain of N fixing bacteria. Perennial legumes fix N during any time of active growth. In annual legumes, N fixation peaks at flowering. With seed formation, it ceases and the nodules fall off the roots. Rhizobia return to the soil environment to await their next encounter with legume roots. These bacteria remain viable in the soil for 3 to 5 years, but often at too low a level to provide significant optimal N-fixation capacity when legume is replanted (Sustainable Agriculture Network, 1998).

The effectiveness of various types of cover crops or combinations of cover crop species on soil N availability and the productivity of succeeding crops has been extensively evaluated (Smith et al., 1987; Holderbaum et al., 1990; Kuo et al., 1996; Clark et al., 1997; Kuo and Jellum, 2002).

In addition to fixing N, cover crops have been reported to reduce the potential for NO_3^- leaching from farm fields (Owens, 1990; Brandi-Dohrn et al., 1997; Staver and Brinsfield, 1998). In studies reviewed by Meisinger et al. (1991), cover crops reduced both the mass of N leached and NO_3^- concentration of leachate by 20 to 80% compared with no cover crop control. They also determined that grasses and brassicas were two to three times as effective as legumes in reducing NO_3^- leaching. Francis et al. (1998), Shepherd (1999), and Rasse et al. (2000) reported that incorporating a nonleguminous cover crop in a cropping system has reduced NO_3^- leaching because the cover crop can reduce water percolation and also effectively use NO_3^- that would otherwise leach.

Cover crops accumulate inorganic soil N between main crop seasons, and holding it in an organic form, prevent it from leaching. The N is subsequently released to the next crop as the cover crop residue decomposes (Dinnes et al., 2002). Rye (*Secale cereale* L.), a cereal, is recognized as having great potential as a scavenger of residual inorganic N present after corn harvest (Staver and Brinsfield, 1990; Ditch and Alley, 1991; McCracken et al., 1994). McCracken et al. (1994) reported that rye was much more effective than vetch (*Vicia villosa* Roth) in reducing NO_3^- leaching. Furthermore, effective erosion control (Kessavalou and Walters, 1997), reduced soil compaction (Raper et al., 2000b), and suppressed weed emergence (Blum et al., 1997) by use of rye in field cropping systems have been reported. The higher capacity of rye in scavenging residual N compared to vetch was associated with its more rapid establishment of an extensive root system (McCracken et al., 1994).

Increasing Organic Matter Content

The importance of soil organic matter in improving soil fertility and productivity is well known (Allison, 1973; Bauer and Black, 1994; Wilhelm et al., 2004). Soil organic matter stabilizes soil aggregates, makes soil easier to cultivate, increases aeration, and increases soil water holding and buffering capacities; soil organic matter breakdown releases available

nutrients to plants (Carter and Stewart, 1996). The soil organic matter content depends on soil type (Schimel et al., 1994), frequency and type of cultivation (Heenan et al., 1995), cropping and residue management (Grace et al., 1995; Webb et al., 2003), and fertilizer N input (Bhogal et al., 1997).

Soil structural degradation is common in intensively cultivated ecosystems due to the depletion of soil organic matter (Grandy et al., 2002). Elliott (1986) reported that aggregate stability decreased in cultivated land compared with native grassland, and that decreases in soil organic matter content paralleled the decrease in stability. Organic matter in agricultural soils contributes positively to soil fertility, soil tilth, crop production, and overall soil sustainability (Bauer and Black, 1994; Reeves, 1997). Cover crops supply organic matter to the soil by decomposing their residues. Data in Table 5 show that legume cover crops increased both the soil organic carbon and organic nitrogen relative to fallow and nonlegume cover crops.

Continued crop production potentials of soils are directly related to their organic matter contents (Lal, 1998; Mann et al., 2002). Within limits, crop productivity is positively related to the soil organic matter content (Reicosky and Forcella, 1998). Soil organic matter improves soil physical, chemical, and biological properties and consequently crop yields (Doran et al., 1998; Doran, 2002; Franzluebbers, 2002). The primary physical characteristics influenced by soil organic matter are those associated with soil structure soil aggregation and aggregate stability (Six et al., 1999). Soil organic matter compounds bind the primary soil particles in the aggregate, physically and chemically, and this, in turn, increases the stability of the aggregates and limits their breakdown during the wetting process (Emerson, 1997; Golchin et al., 1995; Lado and Ben-Hur, 2004). In turn, aggregates and their stability have tremendous influences on infiltration of water, soil water holding capacity, and aeration as well as mass bulk density and penetration resistance (Carter, 2002).

Table 5. Soil organic carbon and nitrogen in 0–7.5 cm depth as affected by cover crops in a no-tillage system

Crop species	Organic carbon (g kg ⁻¹)	Organic nitrogen (g kg ⁻¹)
Fallow	7.9b ^a	0.58c
Hairy vetch	9.7a	0.80ab
Crimson clover	8.4b	0.65bc
Subterranean clover	10.0a	0.81a
Common vetch	10.2a	0.63c
Rye	8.7b	0.65bc

^aValues followed by same letter in the same column are not significantly different ($P < 0.05$).

Source: Adapted from Hargrove (1986).

Chemical properties influenced by soil organic matter content are soil pH, nutrient availability and cycling, cation exchange capacity, and buffering capacity (Tisdall et al., 1986). Organic matter can also bind and detoxify potentially toxic cations [aluminum (Al), manganese (Mn)]. Although a large proportion of the total soil N remains physically and chemically protected from microbial degradation in the stable soil organic matter pool, and thus is unavailable for immediate plant uptake, more labile fractions of soil organic matter, which are generally much smaller, remain an important source of N (Jenkinson and Parry, 1989). In addition, cover crop residue buffers soil against the forces of raindrop impact and wind shear. Crop residues on the soil surface influence radiation balance and energy fluxes and reduce the rate of evaporation from the soil (Wilhelm et al., 2004). Furthermore, increasing C and N storage in soil using management practices, such as cover cropping may also help to reduce the deleterious effects of global warming by increased sequestration of atmospheric CO₂ and N (Lal and Kimble, 1997; Paustian et al., 1997; Post and Kwon, 2000; Sainju et al., 2002).

Soil organic carbon is one of the most important terrestrial pools for carbon storage and exchange with atmospheric CO₂ (Follett, 2001). With land cultivation the soil organic carbon is exposed to oxidation and lost as atmospheric CO₂. This process decreases soil organic carbon. Increasing or maintaining soil C, the basic constituent of soil organic matter, is an important objective for the sustainable use of soil resources (Lal and Kimble, 1997). Appropriate soil and crop management practices such as conservation tillage, use of cover crops, and efficient use of nutrients and water can improve or maintain soil organic carbon. In Table 6 data are presented for carbon sequestration under different management practices.

Enriching the Soil with Essential Mineral Nutrients

In addition to N, cover crops supply other essential nutrients to subsequent crops, when their tissues decompose. Legumes explore subsoil nutrient pools and capture available nutrients through their extensive root systems (Gathumbi et al., 2003). The enrichment of soil with essential nutrients varied with cover crop, especially with quantity of dry matter produced and concentration of nutrients in the dry tissues. Biomass production of cover crops is also affected by environmental conditions, soil fertility and crop management practices. Furthermore, the enrichment of soil with nutrients is mostly dependent on rooting depths, because the desired action lies in the relative transfer to the plow layer of nutrients found only deeper in the horizons. Rooting depth is not only determined by type of crop but more importantly by soil depth, presence of plow pans, aeration, soil texture, soil fertility, disease, and presence of toxic elements. Table 7 shows nutrient accumulation by a mucuna (*Mucuna cinerecum*) cover crop grown on a Brazilian Inceptisol.

Table 6. Soil and crop management practices associated with carbon sequestration in the United States

Management practice	Carbon sequestration (MMTC yr ⁻¹) ^a
Crop rotation and cover crops	5.1–15.3
Conservation reserve program (CRP)	8.8–13.3
Conservation tillage	17.8–35.7
Crop residue/biomass management	11–67
Fallow reduction	1.4–2.7
Fertilizer management	6–18
Livestock manure	3.6–9.0
Supplemental irrigation	1.0–3.2
Total	54.7–164.2
Average	109.4

^aMMTC = Million metric tons of carbon.
Source: Adapted from Follett (2001).

Cover crops have been reported to improve P uptake of succeeding crops (Cavigelli and Thien, 2003). The improved P uptake of succeeding cover crops is associated with several mechanisms. Residues of cover crops may convert relatively unavailable native and residual fertilizer P to chemical forms more available to succeeding crops. Alfalfa (*Medicago sativa* L.), red clover (*Trifolium pratense* L.), sweet clover (*Melilotus officinalis* L.), and

Table 7. Dry matter yield and nutrient accumulation in a ninetyone days old mucuna (graybean) green manure crop grown on an inceptisols of central Brazil

Dry matter yield/nutrient	Value ^a
Dry matter (kg ha ⁻¹)	7016
N (kg ha ⁻¹)	185
P (kg ha ⁻¹)	12
K (kg ha ⁻¹)	161
Ca (kg ha ⁻¹)	73
Mg (kg ha ⁻¹)	19
Zn (g ha ⁻¹)	229
Cu (g ha ⁻¹)	170
Mn (g ha ⁻¹)	1668
Fe (g ha ⁻¹)	2528

^aValues are averages of two field experiments conducted for two consecutive years.
Source: Fageria (unpublished data).

lupine (*Lupinus albus* L.) can absorb more P than most other crops from soils testing low in phosphorus (P) (Gardner et al., 1983; Braum and Helmke, 1995). On decomposition, organic P in the cover crop tissues could provide a relatively labile form of P to succeeding crops (Tiessen et al., 1994). The residue decomposing process may improve P availability to succeeding crops by releasing CO₂, which forms H₂CO₃ in the soil solution, resulting in the dissolution of primary P-containing minerals (Sharpley and Smith, 1989). In soils with high P-fixing capacities, organic compounds released during decomposition processes may increase P availability by blocking P-adsorption sites (Easterwood and Sartain, 1990) and complex excess Al in acidic soils.

Improving Soil Structure

Soil structure is defined as the combination or arrangement of primary soil particles into secondary units or peds (SSSA, 1997). Soil structure is important in describing the health of agricultural soils (Fageria, 2002). Soil structure is important for water infiltration, aeration and plant root development. Improvement of soil structure or aggregation by the action of living and decaying cover crop tissue is widely reported (Lynch and Bragg, 1985; Boyle et al., 1989; Haynes et al., 1991; Haynes and Francis, 1993). Karlen et al. (1994) reported a significant soil aggregation improvement in corn planted in a no-till system with crop residues compared with crop residues removed. Miller and Dick (1995) reported that cover crops provide greater root activity and C inputs which improve soil aggregation and maintain higher organic C pools compared with conventionally managed (fallow) soil.

Improvement in soil hydraulic conductivity and infiltration by modifying soil structure, aggregate stability and macropores have been reported by Murphy et al. (1993) and Kumar and Goh (2000) by crop residues retained on the soil surface. Similarly, up to eightfold increase in hydraulic conductivity in zero-tillage stubble retained have been reported over treatments where stubble was removed by burning (Valzano et al., 1997). Cassel et al. (1995) reported that tillage practices that leave crop residues on the soil surface can reduce or eliminate surface crusting, increase infiltration, and reduce surface runoff and soil loss while increasing crop yields. Baumhardt and Lascano (1996) reported significant improvement in infiltration rate with crop residues left on soil compared with bare soil.

Reducing Soil Erosion

Loss of topsoil by wind and water erosion caused by poor soil management is by far the largest single factor contributing to deterioration of soil physical, chemical, and biological properties and to the further decline in productivity

of most crop lands (Pierce and Lal, 1994; Fageria et al., 1997; Dabney et al., 2001). Soil erosion removes the top soil layer, which generally contains large amounts of soil organic matter and immobile nutrients. The loss of such top soil layers ultimately reduces crop production. The magnitude of the effect of erosion on yields also varies among soils, crops, and management practices (Lal, 1987).

When the soil surface is exposed to raindrop impact, the permeability of the soils is reduced by seal formation (Morin et al., 1981; Ben-Hur and Letey, 1989). The seal formation reduces the infiltration rate, thus increasing runoff (Morin et al., 1981), and may increase soil loss (Ben-Hur et al., 1992). Reduction in soil erosion by cover crops is associated with increasing in soil organic matter content which improve soil water infiltration and holding capacity. With more infiltration and less runoff from each rainfall event, soil erosion is significantly reduced. Cover crops growing after soybean increased surface cover, and anchor residue, and reduced rill erosion (Kaspar et al., 2001).

Increasing Soil Biological Activity

Soil biological properties are intimately related to the chemical environment in the soil and are important in controlling soil tilth as soil chemical and physical properties (Brye et al., 2004). Soil microorganisms play a crucial role in maintaining soil quality due to their action in nutrient cycling through the decomposition of organic matter and nutrient storage (Turco et al., 1994). Cover crops may provide favorable environmental conditions (moisture, temperature, availability of carbon) for the proliferation of soil microorganisms. The soil microbial biomass is the living component of the soil that comprises mainly bacteria and fungi, including soil microfauna and algae (Kumar and Goh, 2000). Although it accounts for only 1 to 3% of organic C and 2 to 6% of organic N in soil (Jenkinson, 1987), it plays a key role in soil organic matter and nutrient dynamics by acting as both a sink (during immobilization) and as a source (mineralization) of plant nutrients (Kumar and Goh, 2000). Several workers have reported higher microbial biomass in no-till than in conventionally tilled soils (Linn and Doran, 1984; Buchanan and King, 1993; Angers et al., 1993; Ndiaye et al., 2000; Schutter and Dick, 2001). Furthermore, higher populations of bacteria, actinomycetes, fungi, earthworms, and nematodes have been reported in residue mulch than in incorporated residues (McCalla, 1958). In addition to the residue management practice, residue quality affects microbial population, fewer bacteria, and fungal populations were reported on grass/cereal residues than in legume residues (Kumar and Goh, 2000; Tian et al., 1993).

Cover crops may also exert a significant effect on soil vesicular-arbuscular mycorrhiza (VAM). The VAM fungal infection increases in corn after cover cropping (Boswell et al., 1998; Kabir and Koide, 2000), whereas no effect

or even slight negative effects were shown in small grain cereals (Baltruschat and Dehne, 1989). Cover crops may also influence the degradation potential of the soil for pesticides (Thorup-Kristensen et al., 2003). Bottomley et al. (1999) found increased degradation of the herbicide 2,4-D in both surface and subsoil layers after a rye cover crop compared to no cover crop in a vegetable cropping system.

Conserving Soil Moisture

Conserving soil moisture with cover crop residues is widely reported (Smith et al., 1987; Sustainable Agriculture Network, 1998). Cover crops residues left on soil improve infiltration of rain water and also reduce evaporative losses, resulting in less moisture stress during drought periods. Grass type cover crops such as rye, barley, wheat, and sorghum-sudangrass have been reported to be very effective in soil moisture conservation (Sustainable Agriculture Network, 1998). Gallaher (1977) showed that soil remained wetter and crop yields were higher when rye was left as surface mulch than when aboveground parts of the rye were removed in a conservation tillage system. Daniel et al. (1999) reported that rye had the highest biomass of several cover crop species tested and soil had higher water contents under rye. The greatest differences in water contents between mulched and bare soils can be expected during short dry periods (7–14 days), not longer ones (Smith et al., 1987).

Suppressing Weeds

Cover crops control weeds and consequently reduce use of herbicides. Weed suppression by cover crops can be due to shading, competition for nutrients and water (Creamer et al., 1996b; Martin, 1996; Liebman and Davis, 2000). Cover crops displace weeds while they grow and their residues can further suppress weeds (Hoffman et al., 1996; Smeada and Weller, 1996; Moyer et al., 2000). Weed control is usually best from dense cover crop plantings and further, when cover crops are allowed to grow for the longest time possible (Smeada and Putnam, 1988). Rye residues are among the most effective mulches and have been reported to suppress weed growth for up to 6 weeks after rye desiccation (Putnam et al., 1983). Nelson et al. (1991) reported less weed biomass with an annual clover (*Trifolium incarnatum* L.) than with two perennial clovers [*Trifolium pretense* L., *Trifolium repens* L. (red and white)] used as spring cover crops. Similarly, Ross et al. (2001) reported that seven clover species suppressed weed biomass but varied from species to species and management practices. Brandsaeter and Netland (1999) reported that weed suppression is one of the several benefits achieved by including a cover crop in a cropping system.

Decreasing Disease and Insect Problems

The benefits of cover crops in controlling diseases and insects may be associated with breaking pest cycles by growing cover crops between main crops. Cover crops have been reported to be responsible for creating a favorable environment for beneficial predators and parasitoid insects, which control harmful insects (Sustainable Agriculture Network, 1998). Similarly, preceding cover crops have been reported to control many soil-borne pathogenic fungal diseases and nematodes in succeeding cash crops (Sustainable Agriculture Network, 1998).

Improving Yields of Subsequent Crops

Improvements in soil physical, chemical, and biological environment by cover crops are known to improve yields of subsequent crops. Increases in crop yields vary from crop to crop and agroecological regions. Furthermore, yield increases depend on management of cover crops as well as subsequent crops. Agronomically, significant N yield responses of cereals following grain legumes compared with cereal monoculture are frequently measured (Chalk, 1998). The positive N response of the subsequently grown cereal has been attributed to the transfer of biologically fixed N, to N sparing under the antecedent legume, and to less immobilization of nitrate during the decomposition of legume residues (Chalk, 1998). Higher corn yield was achieved following white clover (*Trifolium repens* L.) (7.2 Mg ha^{-1}) and red clover (*Trifolium pratense* L.) (6.7 Mg ha^{-1}) cover crops than following no cover crops (5.7 Mg ha^{-1}) (Hively and Cox, 2001). Peoples and Herridge (1990) reported that responses in grain yield of cereals to previous crops of tropical grain legumes varied from $+0.20$ to $+3.68 \text{ t ha}^{-1}$ compared with cereal-cereal monocrop yields with relative increases being in the range 16 to 353%. Improvement in cotton yield in conservation tillage cropping systems has been reported (Bauer and Busscher, 1996; Raper et al., 2000a; Bauer and Roof, 2004).

Several studies have verified that the amount of N in the soil is the key factor in the response of cereals following legumes compared with cereals following nonlegumes (Rowland et al., 1988; Evans et al., 1991; Chalk et al., 1993). However, the response in grain yield may not be entirely due to the amount of available soil N. Improvement in soil structure, the breaking of insects and diseases cycles which afflict cereal monoculture, and phytotoxic and allelopathic effects of different crop residues have all been implicated in the yield response (Chalk, 1998; Peoples and Herridge, 1990). In Table 8 data are presented to show influence of cover crops on yields of corn, sorghum, and cotton after fallow and cover crops. These data indicate that the legumes cover crops provide yield-influencing benefits compared with a fallow/cash crop cropping system.

DISADVANTAGES OF COVER CROPS

There are no known major disadvantages of cover crops. However, concern has been raised over potential harmful effects of cover crops to the succeeding crop (Karlen and Doran, 1991; Johnson et al., 1998). Cover crop may create N deficiency for the next crop if too much N is immobilized and not released in a timely manner (Vyn et al., 1999). Karlen and Doran (1991) showed that cover crops before corn created an early season N deficiency, and even additional N fertilizer did not make up the difference. Similarly, Martinez and Guiraud (1990), Francis et al. (1998), and Wyland et al. (1995) reported that high C/N ratio cover crops may reduce yield of succeeding cash crops because of N immobilization. It is often stated that net N immobilization is likely to occur following addition of plant material with a C/N ratio above 25 (Paul and Clark, 1989).

Rye has been used successfully as a cover crop in the U.S.-northern corn and soybean belt (Dinnes et al., 2002). However, rye should not be grown to maturity as a cover crop because it can reduce the yield of subsequent crops by

Table 8. Yield of corn, sorghum and cotton followed by cover crops. Values are averages across several nitrogen rates and conventional and no-tillage planting systems

Cropping system	Yield of cash crop (Mg ha ⁻¹)	References
Fallow/corn	5.3	Utomo (1986)
Hairy vetch/corn	7.2	Utomo (1986)
Fallow/corn	3.0	Herbek et al. (1987)
Hairy vetch/corn	4.5	Herbek et al. (1987)
Bigflower vetch/corn	4.5	Herbek et al. (1987)
Fallow/corn	5.3	Ebelhar et al. (1984)
Hairy vetch/corn	7.5	Ebelhar et al. (1984)
Bigflower vetch/corn	5.8	Ebelhar et al. (1984)
Crimson clover/corn	5.8	Ebelhar et al. (1984)
Fallow/corn	5.8	Mitchell and Teel (1977)
Rye/hairy vetch/corn	7.1	Mitchell and Teel (1977)
Rye/corn	5.7	Mitchell and Teel (1977)
Fallow/corn	3.7	Adams et al. (1970)
Hairy vetch/corn	5.5	Adams et al. (1970)
Fallow/sorghum	3.4	Hargrove (1986)
Hairy vetch/sorghum	3.9	Hargrove (1986)
Common vetch/sorghum	3.8	Hargrove (1986)
Crimson clover/sorghum	4.1	Hargrove (1986)
Subterranean clover/sorghum	3.9	Hargrove (1986)
Fallow/cotton	0.72	Touchton et al. (1984)
Common vetch/cotton	0.88	Touchton et al. (1984)
Crimson clover/cotton	0.81	Touchton et al. (1984)

using too much water in the spring or immobilizing large amounts of soil N (Tollenaar et al., 1993). Thelen et al. (2004) reported that moisture stress from the interseeded rye was a predominate factor in soybean grain yield reduction. Other studies have shown that a rye cover crop may reduce subsequent corn yields because of allelopathic effects (Raimbault et al., 1990), or N immobilization under low N conditions (Ebelhar et al., 1984; Blevins et al., 1990). Further research may identify rye genotypes that do not release these compounds (Dinnes et al., 2002).

Overall, growing a cover crop rarely causes pest problems. But certain cover crops occasionally may contribute to particular pest, disease, or nematode problems in localized areas, for example by serving as an alternate host to the pest (Sustainable Agriculture Network, 1998). These negative effects of cover crops apply to specific conditions and can be balanced against positive effects that are discussed in the preceding sections.

CONCLUSIONS

In recent years the importance of cover crops in crop production is increasing due to concern for improving soil quality and reducing chemical inputs. Cover crops can provide numerous benefits related to improving soil fertility, structure, water retention, and groundwater quality and reducing soil erosion, and improving pest management. Selecting the right cover crop is critical to successful cover cropping. Furthermore, as with cash crops, cover crops should be rotated periodically to avoid the build-up of plant-specific pests. With proper selection, use, and management of cover crops, it is possible to improve productivity and also could contribute to improved soil, water, and environmental quality.

REFERENCES

- Adams, W.E., Morris, H.D., and Dawson, R.N. (1970) Effect of cropping systems and nitrogen levels on corn (*Zea mays*) yields in the southern Piedmont region. *Agronomy Journal*, 62: 655–659.
- Allison, F.E. (1973) *Soil Organic Matter and its Role in Crop Production*; Elsevier Scientific: New York.
- Angers, D.A., Bissonette, N., and Legere, A. (1993) Microbial and biochemical changes induced by rotation and tillage in a soil under barley production. *Canadian Journal of Soil Science*, 73: 39–50.
- Baltruschat, H. and Dehne, H.W. (1989) The occurrence of vesicular-arbuscular mycorrhiza in agro-ecosystems. II. Influence of nitrogen fertilization and green manure in continuous monoculture and in the crop rotation on the inoculum potential of winter barley. *Plant and Soil*, 113: 251–256.
- Bauer, A. and Black, A.L. (1994) Quantification of the effect of organic matter content on soil productivity. *Soil Science Society of America Journal*, 58: 185–193.

- Bauer, P.J. and Busscher, W.J. (1996) Winter cover and tillage influences on coastal plain cotton production. *Journal of Production Agriculture*, 9: 50–54.
- Bauer, P.J. and Reeves, D.W. (1999) A comparison of winter cereal species and planting dates as residue cover for cotton grown with conservation tillage. *Crop Science*, 39: 1824–1830.
- Bauer, P.J. and Roof, M.E. (2004) Nitrogen, aldicarb, and cover crop effects on cotton yield and fiber properties. *Agronomy Journal*, 96: 369–376.
- Baumhardt, R.L. and Lascano, R.J. (1996) Rain infiltration as affected by wheat residue amount and distribution in ridged tillage. *Soil Science Society of America Journal*, 60: 1908–1913.
- Ben-Hur, M. and Letey, J. (1989) Effect of polysaccharides, clay dispersion, and impact energy on water infiltration. *Soil Science Society of America Journal*, 53: 233–238.
- Ben-Hur, M., Stern, R., van der Merwe, A.J., and Shainberg, I. (1992) Slope and gypsum effects on infiltration and erodibility of dispersive and nondispersive soils. *Soil Science Society of America Journal*, 56: 1571–1576.
- Bhogal, A., Young, S.D., Sylvester-Bradley, R., O'Donnell, F.M., and Ralf, R.N. (1997) Cumulative effects of nitrogen application to winter wheat at Ropsley, U. K. from 1978 to 1990. *Journal of Agricultural Science*, 129: 1–12.
- Blevins, P.L. and Frye, W.W. (1993) Conservation tillage: An ecological approach to soil management. *Advances in Agronomy*, 51: 34–73.
- Blevins, R.L., Herbek, J.H., and Frye, W.W. (1990) Legume cover crops as a nitrogen source for no-till corn and grain sorghum. *Agronomy Journal*, 82: 769–772.
- Blum, U., King, L.D., Gerig, T.M., Lehman, M.E., and Worsham, A.D. (1997) Effects of clover and grain crops and tillage techniques on seedling emergence of some dicotyledonous weed species. *American Journal of Alternative Agriculture*, 12: 146–161.
- Boswell, E.P., Koide, R.T., Shumway, D.L., and Addy, H.D. (1998) Winter wheat cover cropping. A mycorrhizal fungi and maize growth and yield. *Agriculture Ecosystems and Environment*, 67: 55–65.
- Bottomley, P.T., Sawyer, T.E., Boersma, L., Dick, R.P., and Hemphill, D.D. (1999) Winter cover crop enhances 2,4-D mineralization potential of surface and subsurface soil. *Soil Biology and Biochemistry*, 31: 849–857.
- Boyle, M., Frankenberger, W.T., Jr., and Stolzy, L.H. (1989) The influence of organic matter on soil aggregation and water infiltration. *Journal of Production Agriculture*, 2: 290–299.
- Brady, N.C. and Weil, R.R. (2002) *The Nature and Properties of Soils*, 13th ed.; Prentice Hall: Upper Saddle River, New Jersey, 960.
- Brandi-Dohrn, F.M., Dick, R.P., Hess, M., Kauffman, S.M., Hemphill, D.D., and Selker, J.S. (1997) Nitrate leaching under a cereal rye cover crop. *Journal of Environmental Quality*, 26: 181–188.
- Brandsaeter, L.O. and Netland, J. (1999) Winter annual legumes for use as cover crops in row crops in northern regions. I. Field experiments. *Crop Science*, 39: 1369–1379.
- Braum, S.M. and Helmke, P.A. (1995) White lupin utilizes soil phosphorus that is unavailable to soybean. *Plant and Soil*, 176: 95–100.
- Brye, K.R., Slaton, N.A., Mozaffari, M., Savin, M.C., Norman, R.J., and Miller, D.M. (2004) Short-term effects of land leveling on soil chemical properties and their relationships with microbial biomass. *Soil Science Society of America Journal*, 68: 924–934.

- Buchanan, M. and King, L.D. (1993) Carbon and phosphorus losses from decomposing crop residues in no-till and conventional till agroecosystems. *Agronomy Journal*, 85: 631–638.
- Carter, M.R. (2002) Soil quality for sustainable land management: Organic matter and aggregation interactions that maintain soil function. *Agronomy Journal*, 94: 38–47.
- Carter, M.R. and Stewart, B. (1996) *Structure and Organic Matter Storage in Agriculture Soils*; CRC Press: Boca Raton, Florida.
- Cassel, D.K., Raczkowski, C.W., and Denton, H.P. (1995) Tillage effects on corn production and soil physical conditions. *Soil Science Society of America Journal*, 59: 1436–1443.
- Cavigelli, M.A. and Thien, S.J. (2003) Phosphorus bioavailability following incorporation of green manure crops. *Soil Science Society of America Journal*, 67: 1186–1194.
- Chalk, P.M. (1998) Dynamics of biologically fixed N in legume-cereal rotations: A review. *Australian Journal of Agricultural Research*, 49: 303–316.
- Chalk, P.M., Smith, C.J., Hamilton, S.D., and Hopmans, P. (1993) Characterization of the N benefit of a grain legume (*Lupinus angustifolius* L.) to a cereal (*Hordeum vulgare* L.) by an in situ ^{15}N isotope dilution technique. *Biology and Fertility of Soils*, 15: 39–44.
- Chapman, A.L. and Myers, R.J.K. (1987) Nitrogen contributions by grain legumes to rice grown in rotation on the cumunurra soils of the ord irrigation area, Western Australia. *Australian Journal of Experimental Agriculture*, 27: 155–163.
- Clark, A.J., Decker, A.M., Meisinger, J.J., and McIntosh, M.S. (1997) Kill date of vetch, rye, and a vetch-rye mixture. I. Cover crop and corn nitrogen. *Agronomy Journal*, 89: 427–434.
- Creamer, N.G. and Baldwin, K.R. (2000) An evaluation of summer cover crops for use in vegetable production systems in North Carolina. *HortScience*, 35: 600–603.
- Creamer, N.G. and Dabney, S.M. (2002) Killing cover crops mechanically: Review of recent literature and assessment of new research results. *American Journal of Alternative Agriculture*, 17: 32–40.
- Creamer, N.G., Bennett, M.A., Stinner, B.R., and Cardina, J. (1996a) A comparison of four processing tomato production systems differing in cover crop and chemical inputs. *Journal American Society of Horticulture Science*, 121: 559–568.
- Creamer, N.G., Bennett, M.A., Stinner, B.R., Cardina, J., and Regnier, E.E. (1996b) Mechanisms of weed suppression in cover crop-based production systems. *HortScience*, 31: 410–413.
- Dabney, S.M., Delgado, J.A., and Reeves, D.W. (2001) Using winter cover crops to improve soil and water quality. *Communications in Soil Science and Plant Analysis*, 32: 1221–1250.
- Daniel, J.B., Abaye, A.O., Alley, M.M., Adcock, C.W., and Maitland, J.C. (1999) Winter annual cover crops in a Virginia no-till cotton production system. II. Cover crop and tillage effects on soil moisture, cotton yield, and cotton quality. *Journal of Cotton Science*, 3: 84–91.
- Dinnes, D.L., Karlen, D.L., Jaynes, D.B., Kaspar, T.C., Hatfield, J.L., Colvin, T.S., and Cambardella, C.A. (2002) Nitrogen management strategies to reduce nitrate leaching in tile-drained Midwestern soils. *Agronomy Journal*, 94: 153–171.
- Ditch, D.C. and Alley, M.M. (1991) Nonleguminous cover crop management for residual N recovery and subsequent crop yields. *Journal of Fertilizer Issues*, 8: 6–13.
- Doran, J.W. (2000) Soil health and global sustainability: Translating science into practice. *Agriculture Ecosystems and Environment*, 88: 119–127.

- Doran, J.W., Elliot, E.T., and Paustian, K. (1998) Soil microbial pools as related to fallow tillage management. *Soil Tillage Research*, 49: 3–18.
- Easterwood, G.W. and Sartain, J.B. (1990) Clover residue effectiveness in reducing orthophosphate sorption on ferric hydroxide coated soil. *Soil Science Society of America Journal*, 54: 1345–1350.
- Ebelhar, S.A., Frye, W.W., and Blevins, R.L. (1984) Nitrogen from legume cover crops for no-tillage corn. *Agronomy Journal*, 76: 51–55.
- Elliott, E.T. (1986) Aggregate structure and carbon, nitrogen, and phosphorus in native cultivated soils. *Soil Science Society of America Journal*, 50: 627–633.
- Emerson, W.W. (1997) Physical properties and structure. In *Soil Factors in Crop Production in a Semi-Arid Environment*; Russell, I.S. and Greacen, E.L., eds.; University of Queensland Press: St. Lucia, QLD, Australia, 78–104.
- Evans, J., Fettell, N.A., Coventry, D.R., O'Connor, G.E., Walscott, D.N., Mahoney, J., and Armstrong, E.L. (1991) Wheat response after temperate crop legumes in south-eastern Australia. *Australian Journal of Agricultural Research*, 42: 31–43.
- Fageria, N.K. (2002) Soil quality versus environmentally based agricultural management practices. *Communications in Soil Science and Plant Analysis*, 33: 2301–2329.
- Fageria, N.K., Baligar, V.C., and Jones, C.A. (1997) *Growth and Mineral Nutrition of Field Crops*, 2nd Ed.; Marcel Dekker, Inc.: New York, 624.
- Follett, R.F. (2001) Soil management concepts and carbon sequestration in cropland soils. *Soil Tillage Research*, 61: 77–92.
- Francis, G.S., Bartley, K.M., and Tabley, F.J. (1998) The effect of winter cover crop management on nitrate leaching losses and crop growth. *Journal of Agricultural Science*, 131: 299–308.
- Franzluebbers, A.J. (2002) Water infiltration and soil structure related to organic matter and its stratification with depth. *Soil Tillage Research*, 66: 197–205.
- Frye, W.W., Blevins, R.L., Smith, M.S., Corak, S.J., and Varco, J.J.. Role of annual legume cover crops in efficient use of water and nitrogen. In *Cropping Strategies for Efficient Use of Water and Nitrogen*; Hargrove, W.L. ed.; American Society of Agronomy: Madison, Wisconsin, Spec. Publ. No. 51, 129–154.
- Gallaher, R.N. (1977) Soil moisture conservation and yield of crops no-till planted in rye. *Soil Science Society of America*, 41: 145–147.
- Gardner, W.K., Parbery, D.G., and Barber, D.A. (1983) The acquisition of phosphorus by *Lupinus albus* L. I. Some characteristics of the soil/root interface. *Plant and Soil*, 70: 391–402.
- Gathumbi, S.M., Cadisch, G., Buresh, R.J., and Giller, K.E. (2003) Subsoil nitrogen capture in mixed legume stands as assessed by deep nitrogen-15 placement. *Soil Science Society of America Journal*, 67: 573–582.
- Golchin, A., Clarke, P., Oades, J.M., and Skjemstad, J.O. (1995) The effects of cultivation on the composition of organic matter and structural stability of soils. *Australian Journal of Soil Research*, 33: 975–993.
- Grace, P.R., Oades, J.M., Keith, H., and Hancock, T.W. (1995) Trends in wheat yields and soil organic carbon in the permanent rotation trial at the Waite Agricultural Research Institute, South Australia. *Australian Journal of Experimental Agriculture*, 35: 857–864.
- Grandy, A.S., Porter, G.A., and Erich, M.S. (2002) Organic amendment and rotation crop effects on the recovery of soil organic matter and aggregation in potato cropping systems. *Soil Science Society of America Journal*, 66: 1311–1319.
- Hargrove, W.L. (1986) Winter legumes as a nitrogen source for no-till grain sorghum. *Agronomy Journal*, 78: 70–74.

- Haynes, R.J. and Francis, G.S. (1993) Changes in microbial biomass C, soil carbohydrate composition and aggregate stability induced by growth of selected crop and forage species under field conditions. *Journal of Soil Science*, 44: 665–675.
- Haynes, R.J., Swift, R.S., and Stephen, R.C. (1991) Influence of mixed cropping rotations (pastureable) on organic matter content, water stable aggregation and cloud porosity in a group of soils. *Soil Tillage Research*, 19: 77–87.
- Heenan, D.P., McGhie, W.J., Thomson, F.M., and Cham, K.Y. (1995) Decline in soil organic carbon and total nitrogen in relation to tillage, stubble management, and rotations. *Australian Journal of Experimental Agriculture*, 35: 877–884.
- Heichel, G.H. (1987) Legume nitrogen: Symbiotic fixation and recovery by subsequent crops. In *Energy in plant nutrition and pest control*; Helsel, Z.R., ed.; Elsevier Scientific Publishers: Amsterdam, 63–80.
- Herbek, J.H., Frye, W.W., and Blevins, R.L. (1987) Nitrogen from legume cover crops for no-till corn and grain sorghum. In *The Role of Legumes in Conservation Tillage Systems*; Power, J.F., ed.; Soil Conservation Society of America: Ankeny, Iowa, 51–52.
- Hively, W. and Cox, W.J. (2001) Interseeding cover crops into soybean and subsequent corn yields. *Agronomy Journal*, 93: 308–313.
- Hoffman, M.L., Weston, L.A., Snyder, J.C., and Regnier, E.E. (1996) Allelopathic influence of germinating seeds and seedlings of cover crops on weed species. *Weed Science*, 44: 579–584.
- Holderbaum, J.F., Decker, A.M., Meisinger, J.J., Mulford, F.R., and Vough, L.R. (1990) Harvest management of a crimson clover cover crop for no-till corn production. *Agronomy Journal*, 82: 918–923.
- Hoyt, G.D. and Hargrove, W.L. (1986) Legume cover crops for improving crop and soil management in the southern United States. *HortScience*, 21: 397–402.
- Jenkinson, D.S. (1987) Determination of microbial biomass carbon and nitrogen in soil. In *Advances in Nitrogen Cycling in Agricultural Systems*; Wilson, J.R., ed.; CAB International: Wallingford, U.K., 367–386.
- Jenkinson, D.S. and Parry, L.C. (1989) The nitrogen cycle in the Broadbalk wheat experiment: A model for the turnover of nitrogen through microbial biomass. *Soil Biology and Biochemistry*, 21: 535–541.
- Johnson, T.J., Kaspar, T.C., Kohler, K.A., Corak, S.J., and Logsdon, S.D. (1998) Oat and rye overseeded into soybean as fall cover crops in the upper Midwest. *Journal of Soil and Water Conservation*, 53: 276–279.
- Kabir, Z. and Koide, R.T. (2000) The effect of dandelion or a cover crop on mycorrhiza inoculum potential, soil aggregation and yield of maize. *Agriculture Ecosystems and Environment*, 78: 167–174.
- Karlen, D.L. and Doran, J.W. (1991) Cover crop management effects on soybean and corn growth in an on-farm study. *American Journal of Alternative Agriculture*, 6: 71–81.
- Karlen, D.L., Wollenhaupt, N.C., Erbach, D.C., Berry, E.C., Swan, J.B., Eash, N.S., and Jordahl, J.L. (1994) Crop residue effects on soil quality following 10-years of no-till corn. *Soil Tillage Research*, 31: 149–167.
- Kaspar, T.C., Radke, J.K., and Lafen, J.M. (2001) Small grain cover crops and wheel traffic effects on infiltration, runoff, and erosion. *Journal of Soil and Water Conservation*, 56: 160–164.
- Kessavalou, A. and Walters, D.T. (1997) Winter rye as a cover crop following soybean under conservation tillage. *Agronomy Journal*, 89: 68–74.

- Kumar, K. and Goh, K.M. (2000) Crop residues and management practices: Effects on soil quality, soil nitrogen dynamics, crop yield, and nitrogen recovery. *Advances in Agronomy*, 68: 197–319.
- Kuo, S. and Jellum, E.J. (2002) Influence of winter cover crop and residue management on soil nitrogen availability and corn. *Agronomy Journal*, 94: 501–508.
- Kuo, S., Sainju, U.M., and Jellum, E.J. (1996) Winter cover cropping influence on nitrogen mineralization, presidedress soil nitrate test, and corn yields. *Biology and Fertility of Soils*, 22: 310–317.
- Lado, M. and Ben-Hui, M. (2004) Organic matter and aggregate size interactions in infiltration, seal formation and soil loss. *Soil Science Society of America Journal*, 68: 935–942.
- Lal, R. (1987) Effect of soil erosion on crop productivity. *CRC Critical Reviews in Plant Science*, 5: 303–367.
- Lal, R. (1998) Soil erosion impact on agronomic productivity and environmental quality. *Critical Reviews in Plant Science*, 17: 319–464.
- Lal, R. and Kimble, J.M. (1997) Conservation tillage for carbon sequestration. *Nutrient Cycling in Agroecosystems*, 49: 243–253.
- Langdale, G.W., Blevins, R.L., Karlen, D.L., McCool, D.K., Nearing, M.A., Skidmore, E.I., Thomas, A.W., Tyler, D.D., and Williams, J.R.. Cover crop effects on soil erosion by wind and water. In *Cover Crop for Clean Water*; Hargrove, W.L., ed.; Soil and Water Conservation Society: Ankeny, Iowa, 15–23.
- Liebman, M. and Davis, A.S. (2000) Integration of soil, crop and weed management in low-external input farming systems. *Weed Research*, 40: 27–47.
- Linn, D.M. and Doran, J.W. (1984) Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and nontilled soils. *Soil Science Society of America Journal*, 48: 1267–1272.
- Lynch, J.M. and Bragg, E. (1985) Microorganisms and soil aggregate stability. *Advances in Soil Science*, 2: 133–171.
- Mann, L., Tolbert, V., and Cushman, J. (2002) Potential environmental effects on interrow runoff and infiltration. *Soil Science Society of America Journal*, 89: 149–166.
- Martin, C.D. (1996) Weed control in tropical ley farming systems: A review. *Australian Journal of Experimental Agriculture*, 36: 1013–1023.
- Martinez, J. and Guiraud, G.A. (1990) Lysimeter study of the effects of a ryegrass catch crop during a winter wheat/maize rotation, on nitrate leaching and on the following crop. *Journal of Soil Science*, 41: 5–16.
- McCalla, T.M. (1958) Microbial and related studies of stubble mulching. *Journal of Soil and Water Conservation*, 13: 255–258.
- McCracken, D.V., Smith, M.S., Grove, J.H., MacKown, C.T., and Blevins, R.L. (1994) Nitrate leaching as influenced by cover cropping and nitrogen source. *Soil Science Society of America Journal*, 58: 1476–1483.
- Meisinger, J.J., Hargrove, W.L., Mikkelsen, R.L., Williams, J.R., and Benson, V.W.. Effects of cover crops on groundwater quality. In *Cover Crop for Clean Water*; Hargrove, W.L., ed.; Soil and Water Conservation Society: Ankeny, Iowa, 9–11.
- Miller, M. and Dick, R.P. (1995) Dynamics of soil C and microbial biomass in whole soil and aggregates in two cropping systems. *Applied Soil Ecology*, 2: 253–261.
- Mitchell, W.H. and Teel, M.R. (1977) Winter annual cover crops for no-tillage corn production. *Agronomy Journal*, 69: 569–573.
- Morin, J., Benyamini, Y., and Michaeli, A. (1981) The effect of raindrop impact on the dynamics of soil surface crusting and water movement in the profile. *Journal of Hydrology*, 52: 321–335.

- Morse, R.D. (1993) Components of sustainable production systems for vegetables conserving soil moisture. *Horticulture Technology*, 3: 211–214.
- Moyer, J.R., Blackshaw, R.E., Smith, E.G., and McGinn, S.M. (2000) Cereal cover crops for weed suppression in a summer fallow-wheat cropping sequence. *Canadian Journal of Plant Science*, 80: 441–449.
- Murphy, B.W., Koen, T.B., Jones, A., and Huxedrup, L.M. (1993) Temporal variation of hydraulic properties for some soils with fragile structure. *Australian Journal of Soil Research*, 31: 179–197.
- Ndiaye, E.I., Sandeno, J.M., McGrath, D., and Dick, R.P. (2000) Integrative biological indicators for detecting changes in soil quality. *American Journal of Alternative Agriculture*, 15: 26–36.
- Nelson, W.A., Kahn, B.A., and Roberts, B.W. (1991) Screening cover crops for use in conservation tillage systems for vegetables following spring plowing. *HortScience*, 26: 860–862.
- Owens, I.B. (1990) Nitrate-nitrogen concentrations in percolate from lysimeters planted to a legume-grass mixture. *Journal of Environmental Quality*, 19: 131–135.
- Oyer, L.J. and Touchton, J.T. (1990) Utilization legume cropping systems to reduce nitrogen fertilizer requirements for conservation-tilled corn. *Agronomy Journal*, 82: 1123–1127.
- Paul, E.A. and Clark, F.E. (1989) *Soil Microbiology and Biochemistry*; Academic Press.: Toronto, ON, Canada.
- Paustian, K., Andren, O., Janzen, H.H., Lal, R., Smith, P., Tian, G., Tiessen, H., Noordwijk, M.V., and Woomer, P.L. (1997) Agricultural soils as a sink to mitigate CO₂ emissions. *Soil Use and Management*, 13: 230–244.
- Peoples, M.B. and Herridge, D.F. (1990) Nitrogen fixation by legumes in tropical and subtropical agriculture. *Advances in Agronomy*, 44: 155–223.
- Pierce, F.J. and Lal, R. (1994) Monitoring the impact of soil erosion on crop productivity. In *Soil Erosion Research Methods*; Lal, R., ed.; St. Lucie Press: Delray Beach, Florida, 235–263.
- Post, W.M. and Kwon, K.C. (2000) Soil carbon sequestration and landuse change: Processes and potential. *Global Change Biology*, 6: 317–328.
- Putnam, A.R., DeFrank, J., and Barnes, J.P. (1983) Exploitation of allelopathy for weed control in annual and perennial cropping systems. *Journal of Chemical Ecology*, 9: 1001–1010.
- Raimbault, B.A., Vyn, T.J., and Tollenaar, M. (1990) Corn response to rye cover crop management and spring tillage systems. *Agronomy Journal*, 82: 1088–1093.
- Ranells, N.N. and Waggoner, M.G. (1996) Nitrogen release from grass and legume cover crop monocultures and bicultures. *Agronomy Journal*, 88: 777–782.
- Raper, R.L., Reeves, D.W., Burmester, C.H., and Schwab, E.B. (2000a) Tillage depth, tillage timing, and cover crop effects on cotton yield, soil strength, and tillage energy requirement. *Applied Engineering in Agriculture*, 16: 379–385.
- Raper, R.L., Reeves, D.W., Schwab, E.B., and Burmester, C.H. (2000b) Reducing soil compaction of Tennessee Valley soils in conservation tillage systems. *Journal of Cotton Science*, 4: 84–90.
- Rasse, D.P., Ritchie, J.T., Peterson, W.R., Weil, J., and Smucker, A.J.M. (2000) Rye cover crop and nitrogen fertilization effects on nitrate leaching in inbred maize fields. *Journal of Environmental Quality*, 29: 298–304.
- Reeves, D.W. (1994) Cover crops and rotations. In *Crops Residue Management*; Hatfield, J.T. and Stewart, B.A., eds.; Lewis Publishers: Boca Rotan, Florida, 125–172.

- Reeves, D.W. (1997) The role of organic matter in maintaining soil quality in continuous cropping systems. *Soil Tillage Research*, 43: 131–167.
- Reeves, D.W., Wood, C.W., and Touchton, J.T. (1993) Timing nitrogen applications for corn in a winter legume conservation-tillage system. *Agronomy Journal*, 85: 98–106.
- Reicosky, D.C. and Forcella, F. (1998) Cover crop and soil quality interaction in agroecosystems. *Journal of Soil and Water Conservation*, 53: 224–229.
- Ristaino, J.B., Parra, G., and Campbell, C.L. (1996) Suppression of *Phytophthora* blight in bell pepper by a no-till wheat cover crop. *Phytopathology*, 87: 242–249.
- Robertson, G.P., Paul, E.A., and Harwood, R.R. (2000) Greenhouse gases in intensive agriculture: Contributions of individual gases to the radiative forcing of the atmosphere. *Science*, 289: 1922–1925.
- Ross, S.M., King, J.R., Izairralde, R.C., and O'Donovan, J.T. (2001) Weed suppression by seven clover species. *Agronomy Journal*, 93: 820–827.
- Rowland, I.C., Mason, M.G., and Hamblin, J. (1988) Effect of lupines and wheat on the yield of subsequent wheat crops grown at several rates of applied nitrogen. *Australian Journal of Experimental Agriculture*, 28: 91–97.
- Ruffo, M.L. and Bollero, G.A. (2003) Modeling rye and hairy vetch residue decomposition as a function of degree days and decomposition days. *Agronomy Journal*, 95: 900–907.
- Sainju, U.M., Singh, B.P., and Yaffa, S. (2002) Soil organic matter and tomato yield following tillage, cover cropping, and nitrogen fertilization. *Agronomy Journal*, 94: 594–602.
- Sarrantonio, M. and Scoot, T.W. (1988) Tillage effects on availability of nitrogen to corn following a winter green manure crop. *Soil Science Society of America Journal*, 52: 1661–1668.
- Schimel, D.S., Braswell, B.H., and Holland, E.A. (1994) Climatic, edaphic and biotic controls over storage and turnover of carbon in soils. *Global Biogeochemical Cycles*, 8: 279–293.
- Schutter, M.E. and Dick, R.P. (2001) Microbial community profiles and activities among aggregates of winter fallow and cover cropped soil. *Soil Science Society of America Journal*, 66: 142–153.
- Sharpley, A.N. and Smith, S.J. (1989) Mineralization and leaching of phosphorus from soil incubated with surface-applied and incorporated crop residue. *Journal of Environmental Quality*, 18: 101–105.
- Shepherd, M.A. (1999) The effectiveness of cover during eight years of a UK sandland rotation. *Soil Use Management*, 15: 41–48.
- Singer, J.W. and Cox, W.J. (1998) Agronomics of corn production under different crop rotations. *Journal of Production Agriculture*, 11: 462–468.
- Singh, Y., Singh, B., and Khind, C.S. (1992) Nutrient transformations in soils amended with green manures. *Advances in Soil Science*, 20: 237–309.
- Singh, Y., Singh, B., Ladha, J.K., Khind, C.S., Gupta, R.K., Meelu, O.P., and Pasuquin, E. (2004) Long-term effects of organic inputs on yield and soil fertility in the rice-wheat rotation. *Soil Science Society of America Journal*, 68: 845–853.
- Six, J., Elliott, E.T., and Paustian, K. (1999) Aggregate and soil organic matter dynamics under conventional and no-tillage systems. *Soil Science Society of America Journal*, 63: 1350–1358.
- Smeda, R.J. and Putnam, A.R. (1988) Cover crop suppression of weeds and influence on strawberry yields. *HortScience*, 23: 132–134.

- Smeda, R.J. and Weller, S.C. (1996) Potential of rye (*Secale cereale*) for weed management in transplant tomatoes (*Lycopersicon esculentum*). *Weed Science*, 44: 596–602.
- Smith, M.S., Frye, W.W., and Varco, J.J. (1987) Legume winter cover crops. *Advances in Soil Science*, 7: 95–139.
- SSSA (1997) *Glossary of soil science term*; Soil Science Society: Madison, Wisconsin, 134.
- Staver, K.W. and Brinsfield, R.B. (1990) Patterns of soil nitrate availability in corn production systems: Implications for reducing groundwater contamination. *Journal of Soil and Water Conservation*, 45: 318–323.
- Staver, K.W. and Brinsfield, R.B. (1998) Using cereal grain winter cover crops to reduce groundwater nitrate contamination in the Mid-Atlantic Coastal Plain. *Journal of Soil and Water Conservation*, 53: 230–240.
- Sustainable Agriculture Network (1998) *Managing Cover Crops Profitably*, 2nd edition; Handbook Series Book 3, Sustainable Agriculture Network: Beltsville, Maryland, 212 pp.
- Sweeney, D.W. and Moyer, J.L. (2004) In-season nitrogen uptake by grain sorghum following legume green manures in conservation tillage systems. *Agronomy Journal*, 96: 510–515.
- Teasdale, J.R. and Shirley, D.W. (1998) Influence of herbicide application timing on corn production in a hairy vetch cover crop. *Journal of Production Agriculture*, 11: 121–125.
- Thelen, K.D., Mutch, D.R., and Martin, T.E. (2004) Utility of interseeded winter cereal rye in organic soybean production systems. *Agronomy Journal*, 96: 281–284.
- Thorup-Kristensen, K., Magid, J., and Jensen, L.S. (2003) Catch crops and green manures as biological tools in nitrogen management in temperate zones. *Advances in Agronomy*, 79: 227–302.
- Tian, G., Brussaard, L., and Kang, B.T. (1993) Biological effects of plant residues with contrasting chemical compositions under humid tropical conditions: Effects on soil fauna. *Soil Biology and Biochemistry*, 25: 731–737.
- Tiessen, H., Steqart, J.W.B., and Oberson, A. (1994) Innovative soil phosphorus availability indices: Assessing organic phosphorus. In *Soil Testing: Prospects for Improving Nutrient Recommendations*; Havlin, J.L. and Jacobsen, J.S., eds.; SSSA and ASA: Madison, Wisconsin, Spec. Publ. No. 40, 143–162.
- Tisdall, J.M., Nelson, W.L., and Beaton, J.D. (1986) *Soil Fertility and Fertilizers*, 4th ed.; Macmillan Publ. Co.: New York.
- Tollenaar, M.M., Mihajlovic, M., and Vyn, T.J. (1993) Corn growth following cover crops: Influence of cereal cultivar, cereal removal and nitrogen rate. *Agronomy Journal*, 85: 251–255.
- Touchton, J.T., Ricker, D.H., Walker, R.H., and Snipes, C.E. (1984) Winter legumes as a nitrogen source for no-tillage cotton. *Soil Tillage Research*, 4: 391–401.
- Triplett, G.B., Dabney, S.M., and Siefker, J.H. (1996) Tillage systems for cotton on silty upland soils. *Agronomy Journal*, 88: 507–512.
- Turco, R.F., Kennedy, A.C., and Jawson, M.D. (1994) Microbial indicators of soil quality. In *Defining Soil Quality for a Sustainable Environment*; Doran, J.W., Coleman, D.C., Bezdiek, D.F. and Stewart, B.A., eds.; SSSA: Madison, Wisconsin, Spec. Publ. No. 35, 73–90.
- Utomo, M. (1986) *Role of Legume Cover Crops in No-Tillage and conventional Tillage Corn Production*, Ph.D. Dissertation. Lexington, Kentucky: University of Kentucky.

- Valzano, F.P., Greene, R.S.B., and Murphy, B.W. (1997) Direct effects of stubble burning on soil hydraulic properties in a direct drill tillage system. *Soil Tillage Research*, 42: 209–219.
- Vyn, T.J., Janovicek, K.L., Miller, M.H., and Beauchamp, E.G. (1999) Soil nitrate accumulation and corn response to preceding small-grain fertilization and cover crops. *Agronomy Journal*, 91: 17–24.
- Wagger, M.G. (1989) Cover crop management and nitrogen rate in relation to growth and yield of no-till corn. *Agronomy Journal*, 81: 533–538.
- Webb, J., Bellamy, P., Loveland, P.J., and Goodlass, G. (2003) Crop residue returns and equilibrium soil organic carbon in England and Wales. *Soil Science Society of America Journal*, 67: 928–936.
- Wilhelm, W.W., Johnson, J.M.F., Hatfield, J.L., Voorhees, W.B., and Linden, D.R. (2004) Crop and soil productivity response to corn residue removal: A literature review. *Agronomy Journal*, 96: 1–17.
- Wyland, L.J., Jackson, L.E., and Schulbach, K.F. (1995) Soil plant nitrogen dynamics following incorporation of a mature rye cover crop in a lettuce production system. *Journal of Agricultural Science*, 124: 17–25.